

# A Panel Approach for Diagnosing Borreliosis?

by Jyotsna S. Shah, Iris Du Cruz, and Steven Harris

IGenex Inc., Palo Alto, California

## References

1. Tappe J, Jordan D, Janneck E, Fingerle V, Strube C. Revisited: *Borrelia burgdorferi* sensu lato infections in hard ticks (*Ixodes ricinus*) in the city of Hanover (Germany). *Parasites & Vectors*. 2014;7:441.
2. Krause PJ, Fish D, Narasimhan S, Barbour AG. *Borrelia miyamotoi* infection in nature and in humans. *Clin Microbiol Infect*. 2015; 21(7):631-639. Raoult D, Roux V. The body louse as a vector of reemerging human diseases. *Clin Infect Dis*. 1999; 29(4):888-911.
3. Cutler SJ, Abdissa A, Trape J-F. New concepts for the old challenges of African relapsing fever borreliosis. *Clin Microbiol Infect*. 2009;15:400-406.
4. Hoch M, Wieser A, Löscher T, et al. Louse-borne relapsing fever (*Borrelia recurrentis*) diagnosed in 15 refugees from northeast Africa: epidemiology and preventive control measures, Bavaria, Germany, July to October 2015. *Euro Surveill*. 2015 Oct 22; 20(42).
5. Trape J-F, Diatta G, Arnathau C, et al. The epidemiology and geographic distribution of relapsing fever Borreliosis in West and North Africa, with a review of the *Ornithodoros erraticus* complex (Acari: Ixodida). Bergström S, ed. *PLoS ONE*. 2013; 8(11):e78473. doi:10.1371/journal.pone.0078473.
6. Krause PJ. *Borrelia* species (Relapsing Fever). <http://antimicrobe.org>Cutler SJ. Relapsing fever Borreliae: a global review. *Clin Lab Med*. 2015 Dec; 35(4):847-65.
7. Fukunaga M, Takahashi Y, Tsuruta Y, Matsushita O, Ralph D, McClelland M, Genetic and phenotypic analysis of *Borrelia miyamotoi* sp. nov., isolated from the ixodid tick *Ixodes persulcatus*, the vector for Lyme disease in Japan. *Int J Syst Bacteriol*. 1995; 45:804-810.
8. Platonov AE, Karan LS, Kolyasnikova NM, et al. Humans infected with relapsing fever spirochete *Borrelia miyamotoi*, Russia. *Emerg Infect Dis*. 2011; 17:1816-1823.
9. Gugliotta JL, Goethert HK, Berardi VP, Telford SR 3<sup>rd</sup>. Meningoencephalitis from *Borrelia miyamotoi* in an immunocompromised patient. *New Engl J Med*. 2013;368:240-245.
10. Shah JS, Du Cruz I, Narciso W, Lo W, Harris NS. Improved Sensitivity of Lyme Disease Western Blots Prepared with a Mixture of *Borrelia Burgdorferi* Strains 297 and B31. *Chronic Dis Int*. 2014;1(2):1-7.
11. Scoles GA, Papero M, Beati L, Fish D. A relapsing fever group spirochete transmitted by *Ixodes scapularis* ticks. *Vector Borne Zoonotic Dis*. 2001; 1:21-34. [PubMed: 12653133]
12. Tsao JI, Wootton JT, Bunikis J, Luna MG, Fish D, Barbour AG. An ecological approach to preventing human infection: vaccinating wild mouse reservoirs intervenes in the Lyme disease cycle. *Proc Natl Acad Sci U S A*. 2004; 101:18159-18164. [PubMed: 15608069]
13. Barbour AG, Bunikis J, Travinsky B, et al. Niche partitioning of *Borrelia burgdorferi* and *Borrelia miyamotoi* in the same tick vector and mammalian reservoir species. *Am J Trop Med Hyg*. 2009; 81:1120-1131.
14. Tokarz R, Jain K, Bennett A, Briese T, Lipkin WI. Assessment of polymicrobial infections in New York State. *Vector Borne Zoonotic Dis*. 2010;10:217-221. [PubMed: 19725770]
15. Hamer SA, Hickling GJ, Keith R, et al. Associations of passerine birds, rabbits, and ticks with *Borrelia miyamotoi* and *Borrelia andersonii* in Michigan, USA. *Parasites & Vectors*. 2012; 5:231. [PubMed: 23057837]
16. Crowder CD, Carolan HE, Rounds MA, et al. Prevalence of *Borrelia miyamotoi* in Ixodes ticks in Europe and the United States. *Emerg Infect Dis*. 2014; 20:1678-1682. [PubMed: 25280366]
17. Padgett K, Bonilla D, Kjemtrup A, et al. Large scale spatial risk and comparative prevalence of *Borrelia miyamotoi* and *Borrelia burgdorferi* sensu lato in *Ixodes pacificus*. *PLoS One*. 2014; 21:9, e110853.
18. Fedorova N, Kleinjan JE, James D, Hui LT, Peeters H, Lane RS. Remarkable diversity of tick or mammalian-associated Borreliae in the metropolitan San Francisco Bay Area, California. *Ticks Tick-borne Dis*. 2014; 5:951-961. [PubMed: 25129859]
19. Salkeld D, Cinkovich S, Nieto NC. Tick-borne pathogens in northwestern California, USA. *Emerg Infect Dis*. 2014; 20:493-494. [PubMed: 24565119].