## An Integrative Medical Approach to Macular Degeneration- Part 2

by Marc Grossman, OD, Lac Townsend Letter #442, April 2020

## **Complete References**

- 1. Snellen EL, et al. Neovascular age-related macular degeneration and its relationship to antioxidant intake. *Acta Ophthalmol Scand*. 2002 Aug;80(4):368-71.
- Richer S, et al. Double-masked, placebo-controlled, randomized trial of lutein and antioxidant supplementation in the intervention of atrophic age-related macular degeneration: the Veterans LAST study (Lutein Antioxidant Supplementation Trial). *Optometry*. April 2004;75(4):216-30.
- Souied EH, et al. Oral docosahexaenoic acid in the prevention of exudative age-related macular degeneration: The Nutritional AMD Treatment 2 Study. *Ophthalmology*. February 2013.



- 4. Landrum JT, et al. A one-year study of the macular pigment: the effect of 140 days of a lutein supplement. *Exp Eye Res.* 1997 Jul;65(1):57-62.
- Bernstein PS, et al. Resonance Raman measurement of macular carotenoids in normal subjects and in age-related macular degeneration patients. *Ophthalmology*. 2002;109(10):1780-1787.
- 6. Kijlstra A, et al. Lutein: more than just a filter for blue light. *Prog Ret Eye Res*. July 2012.
- 7. Studies show there is consistent evidence that high intake of docosahexaenoic acid (DHA; 22:6, n-3), a long-chain omega-3 polyunsaturated fatty acid (PUFA) present in

oily fish, is associated with a reduced risk of neovascular AMD. 1.) Chong EW, et al. Dietary omega-3 fatty acid and fish intake in the primary prevention of age-related macular degeneration: a systematic review and meta-analysis. *Arch Ophthalmol*. 2008;126(6): 826–833. 2.) Augood C, et al. Oily fish consumption, dietary docosahexaenoic acid and eicosapentaenoic acid intakes, and associations with neovascular age-related macular degeneration. *Am J Clin Nutr*. 2008;88(2): 398–406. 3.) Merle B, et al. Dietary omega-3 fatty acids and the risk for age-related maculopathy: the Alienor Study. *Invest Ophthalmol Vis Sci*. 2001;52(8): 6004–6011. 4.) Seddon JM, et al. Dietary fat and risk for advanced age-related macular degeneration. *Arch Ophthalmol*. 2001;119(8): 1191–1199. 5.) Merle BM, et al. High concentrations of plasma n3 fatty acids are associated with decreased risk for late age-related macular degeneration. *J Nutr*. 2013;143(4): 505–511.

- SanGiovanni JP, Chew EY. The role of omega-3 long-chain polyunsaturated fatty acids in health and disease of the retina. *Prog Retin Eye Res.* 2005;24(1): 87–138; Tan JSL, et al. Dietary Fatty Acids and the 10-Year Incidence of Age-Related Macular Degeneration: The Blue Mountain Eye Study. *Arch Ophthalmol.* 2009;127(5):656-665.
- 9. Smith W, Mitchell P, Leeder SR. Dietary fat and fish intake and age-related maculopathy. *Arch Ophthalmol*. 2000;118(3): 401–404.
- 10. Cho E, et al. Prospective study of dietary fat and the risk of age-related macular degeneration. *Am J Clin Nutr*. 2001;73(2): 209–218.
- 11. Delcourt C, et al. Dietary fat and the risk of age-related maculopathy: the POLANUT Study. *European Journal of Clinical Nutrition*. February 2007.
- 12. Millen AE, et al. Vitamin D Status and Early Age-Related Macular Degeneration in Postmenopausal Women. *Arch Ophthalmol.* April 2011.
- 13. Hill DJ. Vitamin D may play key role in preventing macular degeneration. August 27, 2016.
- 14. Piermarocchi S, et al. Carotenoids in Age-Related Maculopathy Italian Study: two-year results of a randomized study. *Eur J Opthalmol*. March-April 2012.
- 15. Maher TJ. Astaxanthin, Continuing Education Module. New Hope Institute of Healing;2000.
- 16. Otsuka T, et al. Protective effects of a dietary carotenoid, astaxanthin, against lightinduced retinal damage. *J Pharmacol Sci.* 2013;123(3):209-18.
- 17. Park JS, et al. Astaxanthin decreased oxidative stress and inflammation and enhanced

immune response in humans. Nutr Metab (Lond). 2010;7:18.

- 18. Akuffo KO, et al. Sustained supplementation and monitored response with differing carotenoid formulations in early age-related macular degeneration. *Eye*. May 2015.
- 19. Bone RA, et al. Macular pigment response to a supplement containing meso-zeaxanthin, lutein and zeaxanthin. Nutr Metab (Lond). 2007.
- 20. Anekonda TS, Adamus G. Resveratrol prevents antibody-induced apoptotic death of retinal cells through upregulation of Firt1 and Ku70. *BMC Res Notes*. December 1, 2008.
- 21. Remsberg CM, et al. Pharmacometrics of pterostilbene: preclinical pharmacokinetics and metabolism, anticancer, anti-inflammatory, antioxidant, and analgesic activity. *Phytother Res.* 2008 Feb;22(2):169-79.
- Perecko T, et al. Structure-efficiency relationship in derivatives of stilbene. Comparison of resveratrol, pinosylvin and pterostilbene. *Neuro Endocrinol Lett.* 2008 Oct;29(5):802-5.
- 23. Currais A, et al. Modulation of p25 and inflammatory pathways by fisetin maintains cognitive function in Alzheimer's disease transgenic mice. *Aging Cell*. 2014 Apr;13(2):379-90.
- 24. Antonetti DA, Klein R, Gardner TW. Diabetic retinopathy. *NEJM*. 2012 Mar 29; 366(13):1227-39.
- 25. Vitseva O, et al. Grape see and skin extracts inhibit platelet function and release of reactive oxygen intermediates. *J Cardiovasc Pharmacol*. 2005 Oct;46(4):445-51.
- 26. Balu M, et al. Age-related oxidative protein damages in central nervous system of rats: modulatory role of grape seed extract. *Int J Dev Neurosci*. 2005 Oct;23(6):501-7.
- 27. Du J, Cullen JJ, Buettner GR. Ascorbic acid: chemistry, biology, and the treatment of cancer. *Biochim Biophys Acta*. 2012 Dec;1826(2):443-57.
- 28. Age-Related Eye Disease Study Research Group. A randomized, placebo-controlled, clinical trial of high-dose supplementation with vitamins C and E, beta carotene, and zinc for age-related macular degeneration and vision loss: AREDS report no. 8. Arch Ophthalmol. 2001 Oct;119(10):1417-36.
- 29. Falsini B, et al. Influence of saffron supplementation on retinal flicker sensitivity in early age-related macular degeneration. *Investigations in Opththalmological and Visual Science*. November 2010.

- Marangoni D, et al. Functional effect of Saffron supplementation and risk genotypes in early age-related macular degeneration: a preliminary report. *Journal of Translational Medicine*. September 2013.
- 31. Alavizadeh SH, Hosseinzadeh H. Bioactivity assessment and toxicity of crocin: a comprehensive review. *Food and Chemical Toxicology*. February 2014.
- 32. Corso L, et al. Saffron reduces ATP-induced retinal cytotoxicity by targeting P2X7 receptors. *Purinergic Signaling*. March 2016.
- 33. Broadhead GK, et al. Dietary modification and supplementation for the treatment of agerelated macular degeneration. *Nutrition Reviews*. July 2015.
- 34. Broadhead GK, et al. Efficacy and Safety of Saffron Supplementation: Current Clinical Findings. *Critical Reviews in Food Science and Nutrition*. April 2015.
- 35. Michael LD. Allen MJ. Nutritional supplementation, electrical stimulation and age related macular degeneration. *J Orthomol Med.* 1993: 8: 168-171. In this 7-year clinical study, 400 eyes with AMD, 78% of the eyes showed from 1-9 lines of improvement in reading of the visual acuity chart; Over 50% improved from 2-9 lines; 2 patients with

retinal vein occlusion and swelling of the macula had dramatic improvement in vision.

36. Reader AL, Halloran, G. Bioelectrical Stimulation in an Integrated Treatment for Macular Degeneration, RP, Glaucoma, CMV, and DR; Fourth Annual Symposium on



Biological Circuits, Oct. 1997, Mankato University, MN. In a two-year study involving 114 patients, researchers investigated effects of microcurrent stimulation on patients with macular degeneration, retinitis pigmentosa and other retina problems. Of 18 patients with AMD, 16 showed improvement. Of 78 patients with retinitis pigmentosa, 62 showed improvement. Of 18 patients with other retinal problems, 16 showed improvement;14 other patients showed no improvement, but their condition stabilized and did not get worse as expected. Two other patients continued to lose vision, but at a slow rate.

- 37. Wallace LB. The Treatment of Macular Degeneration and Other Retinal Diseases Using Bioelectromagnetic Therapy. *Journal of Optometric Phototherapy*. March 1997.
- 38. Paul E. The Treatment of Retinal Disease With MCS and Nutritional Supplementation, International Society for Low-Vision Research and rehabilitation at the Low Vision Congress in Gothenberg, Sweden 2002.
- O'Clock GD, Jarding JB. Electrotherapeutic device/protocol design considerations for visual disease applications. Engineering in Medicine and Biology Society, 2009. EMBC 2009. Annual International Conference of the IEEE. 2009:2133-6.
- 40. Chaikin L, et al. Microcurrent stimulation in the treatment of dry and wet macular degeneration. *Clin Ophthalmol* (Auckland, NZ). December 2015;9(1):2345-53.

- 41. Muth et al. The effect of bilberry nutritional supplementation on night visual acuity and contrast sensitivity. *Altern Med Rev.* 2000 Apr;5(2):164-73.
- 42. Is neuroprotective and has been shown to improve microcirculation. *Mol Nutr Food Res*. 2009 Jul;53(7):869-77; *Clin Chem*. 2011 Nov;57(11):1524-33; *Metabolism*. 2000 Jul;49(7):880-5; *NeuroToxicology* 28 (2007) 93–100.
- 43. Christian P, et al. Vitamin A or Beta-Carotene Supplementation Reduces But Does Not Eliminate Maternal Night Blindness in Nepal. *J. Nutr.* September 1998.
- 44. Lebuisson DA, et al. Presse Med. 1986 Sep 25;15(31):1556-8.
- 45. Fies P, Dienel A. [Ginkgo extract in impaired vision treatment with special extract EGb 761 of impaired vision due to dry senile macular degeneration]. *Wien Med Wochenschr*, 2002. (article in German)
- 46. Arch Phys Med Rehabil. 2000 May;81(5):668-78.
- 47. Induction of vascular endothelial growth factor by 4-hydroxynonenal and its prevention by glutathione precursors in retinal pigment epithelial cells. *European Journal of Pharmacology*. August 9, 2002;
- 48. Sternberg, Davidson, Jones, et al. Invest Ophthalmol Vis Sci. 1993 Dec;34(13):3661-8.
- 49. Deficiency of taurine, an amino acid, has been shown to lead to retinal degeneration and supplementing it has been used with some success to prevent, treat and stabilize retinal changes: *Altern Med Rev.* 1998 Apr;3(2):128-36; *Oftalmol Zh.* 1989;(8):463-5; *Brain Res Rev.* 1991 May-Aug;16(2):151-69; *J Neurosci Res.* 1987;18(4):602-14.
- 50. Mares-Perlman JA, et al. Serum antioxidants and age-related macular degeneration in a population-based case-control study. *Arch Ophthalmol*. 1995 Dec;113(12):1518.
- 51. Stefanova NA, et al. Potential of melatonin for prevention of age-related macular degeneration: experimental study (in Russian), *Advances in Gerontology*. 2013.
- 52. Yi C, et al. Effects of melatonin in age-related macular degeneration. *Ann N Y Acad Sci.* 2005 Dec;1057:384-92.
- 53. Jarrett SG, Boulton ME. Consequences of oxidative stress in age-related macular degeneration. *Mol Aspects Med.* 2012 Aug;33(4):399–417.
- 54. Green tea extract has excellent scavenging reactive oxygen species such as superoxide and the hydroxyl and peroxyl radical. *Chem Res Toxicol*. 2003 Sep;16(9):1155-61; *Addict Biol*. 2002 Jul;7(3):307-14; *Arch Biochem Biophys*. 1999 Feb 1;362(1):79-86;

*Phytother Res.* 2010 Apr;24(4):503-9; *Food Chem Toxicol.* 201 Apr;48(4):1032-9; *Free Radic Biol Med.* 2003 Mar 15;34(6):648-62; *Biophys J.* 2009 Feb;96(3):1026-35.

- 55. Study findings strongly suggested that an appropriate combination acetyl-L-carnitine, n-3 fatty acids, and coenzyme Q10 which affect mitochondrial lipid metabolism may improve and subsequently stabilize visual functions, and it may also improve fundus alterations in patients affected by early AMD. *Ophthalmologica*. 2005 May-Jun;219(3):154-66.
- 56. Feher J, et al. Improvement in visual functions and fundus alterations in early age-related macular degeneration treated with a combination of acetyl-L-carnitine, n-3 fatty acids, and coenzyme Q10. *Ophthalmologica*. May-June 2005.
- 57. There is an association between both low serum selenium levels and current smoking status and the development of age-related macular degeneration. Mayer, et al. *Acta Ophthalmol Scand*. 1998 Feb;76(1):62-7.
- 58. Tsang NC, et al. Serum levels of antioxidants and age-related macular degeneration. *Doc Ophthalmol.* 1992;81(4):387–400.
- 59. The researchers found that daily supplementation with the milk/goji formulation increased the levels of zeaxanthin in the blood and protected from additional drusen formulation or loss of pigmentation. Bucheli P, et al. Goji berry effects on macular characteristics and plasma antioxidant levels. *Optometry and Visual Science*. February 2011.
- 60. Zhuang P, et al. Effect of quercetin on formation of choroidal neovascularization (CNV) in age-related macular degeneration (AMD). *Eye Sci*. March 2011;26(1):23-9.
- 61. Benavente-Garcia O, Castillo J. Update on uses and properties of citrus flavonoids: new findings in anticancer, cardiovascular, and anti-inflammatory activity. *J Agric Food Chem.* Aug 13, 2008;56(15):6185-205.
- 62. Saberi-Karimian M, et al. Vascular endothelial growth factor: An important molecular target of curcumin. *Crit Rev Food Sci Nutr*. August 2017; 30:1-14.
- 63. Park SI, et al. Anti-apoptotic effects of Curcuma longa L. extract and its curcuminoids against blue light-induced cytotoxicity in A2E-laden human retinal pigment epithelial cells. *J Pharm Pharacol*. March 2017;69(3):334-340.
- 64. Zhu W, et al. Effect of curcumin on aging retinal pigment epithelial cells. *Drug Des Devel Ther*. September 2015; 25;9:5337-44.
- 65. Shay J, et al. Molecular Mechanisms and Therapeutic Effects of (-)-Epicatechin and

Other Polyphenols in Cancer, Inflammation, Diabetes, and Neurodegeneration. *Oxid Med Cell Logev*. 2015: 181260.

- 66. <u>https://chriskresser.com/how-too-much-omega-6-and-not-enough-omega-3-is-making-us-sick/</u>
- 67. Liew G, et al. The association of aspirin use with age-related macular degeneration. *JAMA Internal Medicine*. 2013;173(4):258-64.
- 68. Andriessen EM, et al. Gut microbiota influences pathological angiogenesis in obesitydriven choroidal neovascularization. *EMBO Molecular Medicine*. December, 2016.
- 69. Rowan S, et al. Involvement of a gut-retina axis in protection against dietary glycemiainduced age-related macular degeneration. *Proceedings of the National Academy of Science, USA*. May 2017. Researchers have noted that the inability of the body to properly process sugar, leading to advanced glycation (bonding of sugar molecules with proteins and other molecules), contributes to development of macular degeneration. This study concluded that the health of gut microbiota was a factor in the ability of the body to process sugars and that this ability was hampered in a high-glycemia diet.
- 70. Less polyunsaturated fat in one's diet may help protect against age-related macular degeneration (AMD), according to results of the Carotenoids in Age-Related Eye Disease Study. Reference: *Arch Ophthalmol*. 2009 Nov;127(11):1483-93.
- 71. Chiu CJ, et al. Dietary carbohydrate and the progression of age-related macular degeneration: A prospective study from the Age-Related Eye Disease Study. *Am J Clin Nutr.* 2007;86(4):1210-8.
- 72. www.jackkruse.com/quatum-biology-5-coherent-water
- 73. www.ncbi.nlm.nih.govpubmed/259550246
- 74. https://www.jackkruse.com/emf-5-what-are-the-biologic-effects-of-emf/
- 75. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4734149/
- 76. <u>www.jackkruse.com/emf-1</u>

## Return to TownsendLetter.com